

Мадумарова Зарнигора Шухрат кизи
Факультет повышения квалификации,
Подготовка и переподготовка врачей,
старший преподаватель лечебного факультета
Радиология и клиническая
лабораторная диагностика
Андижанский государственный
медицинский институт
Андижан, Узбекистан

Назарова Гульчехра Усмоновна
факультет повышения квалификации
Подготовка и переподготовка врачей,
Заведующая кафедрой
семейной медицины, к.м.н.
Андижанский государственный
медицинский институт
Андижан, Узбекистан

УСОВЕРШЕНСТВОВАНИЕ ДИАГНОСТИКИ ОСТЕОАРТРОЗА ТАЗОБЕДРЕННОГО СУСТАВА С ИСПОЛЬЗОВАНИЕМ УЛЬТРАСОНОГРАФИЧЕСКОГО ИССЛЕДОВАНИЯ

For citation: Z.Sh. Madumarova, G.U. Nazarova. IMPROVING THE DIAGNOSIS OF OSTEOARTHRITIS HIP JOINT OF USING ULTRASONOGRAPHY. Journal of cardiorespiratory research. 2023, vol 1.1, issue 55, pp.271-274.

АННОТАЦИЯ

Данная статья посвящена применению УЗИ в диагностике остеоартроза тазобедренного сустава. В статье обсуждается актуальность применения УЗИ при заболевании остеоартрозом (ОА), ревматоидным артритом (РА), а также дифференциальная оценка воспалительных и дегенеративных процессов на основе ультразвуковой доплерографии. Долгое время простой рентген считался эталонным методом остеоартроза (ОА). В последнее время ультразвуковая диагностика является инновационным методом визуализации этого заболевания. Использование ультразвука очень подробно выделяет различные анатомические структуры и обнаруживает внутри- и внесуставные изменения.

Ключевые слова: коксартроз, коксит, ревматоидный артрит, остеоартроз, УЗИ, тазобедренные суставы, диагностика.

Madumarova Zarnigor Shukhrat kizi
senior teacher at the Department of Medical
Radiology and Clinical Laboratory Diagnostics,
Faculty of Advanced
Training and Retraining of Physicians,
Andijan State Medical Institute,
Andijan city, Uzbekistan

Nazarova Gulchexra Usmonovna
PhD, Head of the Department of Family Medicine,
Faculty of Advanced
Training and Retraining of Physicians
Andijan State Medical Institute
Andijan city, Uzbekistan

IMPROVING THE DIAGNOSIS OF OSTEOARTHRITIS HIP JOINT OF USING ULTRASONOGRAPHY

ANNOTATION

This article is devoted to the application of ultrasonography in the diagnosis of osteoarthritis of the hip joint. The article discusses the relevance of the use of ultrasound in the disease of osteoarthritis (OA), rheumatoid arthritis (RA), as well as the differential assessment of inflammatory and degenerative processes based on ultrasound Doppler sonography. For a long time, a simple X-ray was considered the reference technique of osteoarthritis (OA). Recently, ultrasonography is an innovative method for visualizing this disease. The use of ultrasound highlights the various anatomical structures in great detail and detects intra- and extra-articular changes.

Keywords: Coxarthrosis, coxitis, rheumatoid arthritis, osteoarthritis, ultrasonography, hip joints, diagnostics.

Madumarova Zarnigor Shuhrat qizi
malaka oshirish fakulteti
Vrachlarni tayyorlash va qayta tayyorlash,
tibbiyot kafedrasini katta o'qituvchisi
Radiologiya va klinik laboratoriya diagnostikasi,
Andijon davlat tibbiyot instituti,
O'zbekiston, Andijon
Nazarova Gulchexra Usmonovna
malaka oshirish fakulteti Vrachlarni tayyorlash va
qayta tayyorlash,
Oila tibbiyoti kafedrasini mudiri, t.f.n
Andijon davlat tibbiyot instituti,
O'zbekiston, Andijon

ULTRASONOGRAFIK TEKSHIRISH USULIDAN FOYDALANGAN HOLDA CHANOQ-SON BO'G'IMLARI OSTEOARTRITI DIAGNOSTIKASINI TAKOMILLASHTIRISH

ANNOTATSIYA

Ushbu maqola son qo'shimchasining osteoartritini tashxislashda ultratovush tekshiruvidan foydalanishga bag'ishlangan. Maqolada osteoartrit (OA), revmatoid artrit (RA) da ultratovushdan foydalanishning dolzarbligi, shuningdek, ultratovushli Doppler sonografiyasi asosida yallig'lanish va degenerativ jarayonlarni differentsial baholash muhokama qilinadi. Uzoq vaqt davomida oddiy rentgenografiya osteoartritni (OA) o'rganish uchun mos yozuvlar usuli hisoblanadi. So'nggi paytlarda ultratovush tekshiruvini ushbu kasallik uchun innovatsion tasvirlash usuli hisoblanadi. Ultratovush tekshiruvidan foydalanish turli xil anatomik tuzilmalarni juda batafsil tekshirish va intra- va ekstra-artikulyar o'zgarishlarni aniqlash imkonini beradi.

Kalit so'zlar: Koksartroz, koksit, revmatoid artrit, artroz, ultratovush, son bo'g'imlari, diagnostika.

Introduction

Hip joint disease is one of the urgent problems of modern orthopedics, as it contributes to the rapid development of severe anatomical and functional inferiority of the musculoskeletal system, in particular in the elderly. According to WHO, more than ten percent of the world's population suffers from joint diseases. In terms of the frequency of lesions, the hip joint ranks first (42.7%), the second is the knee (33.3%), the third is the shoulder (10.8%), and the remaining joints account for 13.2% [1].

The differential diagnosis of lesions of the hip joint is difficult due to its deep occurrence and the presence of large muscle masses around the joint. In this regard, it is not possible to assess the external changes in the joint, it is very difficult to perform its puncture and arthroscopy. The differential diagnosis of coxarthrosis and coxitis at an early stage is especially difficult [2]. In this regard, the issue of differential diagnosis of diseases of the hip joint of dystrophic and inflammatory origin is relevant.

The ultrasound method, being non-invasive, reproducible and relatively affordable, can be widely used to assess both the accumulation of fluid inside the joint, in the joint bags, and to assess the thickness of the synovial membrane and erosive changes [3,4]. Modern ultrasound equipment makes it possible to evaluate both superficial periarticular and intra-articular tissues that make up the joint.

Purpose of the study: Improving the Efficiency of Radiation Diagnosis of Inflammatory and Degenerative Diseases of the Hip Joint Based on the Ultrasound Method.

Materials and methods of research.

The total number of patients examined was 138, of which 15 (13.9%) were men and 93 were women (86.1%).

The patients were divided into groups and subgroups. Group I included patients with dystrophic diseases (coxarthrosis stage I-III), a total of 82 people. Group II included 10 patients with coxarthrosis formed as a result of aseptic necrosis of the femoral head. Of these, men 5 (50%) people, women 5 (50%) people. Group III with inflammatory diseases included patients with rheumatoid arthritis, in the amount of 12 people. Of these, 4 men (33.3%) people, women 8 (66.7%) people.


All patients in the period from 2018 - 2020. were treated by an orthopedist in the ASMI clinic, and an ultrasound examination was carried out in the radiology department of the ASMI clinic in Andijan.

Ultrasound examination was carried out on digital multifunctional scanners "Sonoscape S-22" (China) and "Mindray DC3" (China). For the most clear visualization, a study was used with convex and linear sensors, in the frequency range of 7-12 MHz.

Comprehensive ultrasound examination included survey scanning of the hip joint in gray scale mode, examination of the shape and contour of the femoral head, determination of the presence of effusion, measurement of parameters of the joint capsule, thickness of the hypoechoic layer of hyaline cartilage, examination of the state of muscles, ligaments and bones adjacent to the joint.

When evaluating the structures of the hip joint according to the standard method from the anterior approach, the bone landmarks were the upper edge of the acetabulum and the semicircle of the femoral head, where hypoechoic hyaline cartilage, the articular capsule of the hip joint were visualized (Fig. 1).

Fig. 1. Sonogram of the structural elements of the hip



joint is normal. The arrows in the diagram show the measurement of the thickness of the hypoechoic layer of hyaline cartilage. 2 - fibrous capsule.

In patients of all groups, the regional blood flow in the lateral circumflex arteries of the thigh was assessed. The assessment of blood flow was carried out using color Doppler mapping and pulsed wave spectral Doppler, while assessing: peak blood flow velocity (PSV), end-diastolic blood flow velocity (DPV), resistance index (IR).

Research results

When analyzing the revealed sonographic and Dopplerographic diagnostic criteria for the stages of the pathological process, in patients of group I, it was revealed that the thickness of the hypoechoic layer of hyaline cartilage decreases from normal values of 2.4 mm to 1.3 mm as the stages of the disease increase in patients of subgroup IA, to 0.9 mm in subgroup IB and thins to 0.8 mm in subgroup I C. At stage III of coxarthrosis, the thickness of the hypoechoic layer of hyaline cartilage is 0.85 mm, and with coxarthrosis formed as a result of aseptic necrosis of the femoral head, cartilage in the necrosis zone is not visualized. In rheumatoid arthritis, there was an increase in the thickness of the fibrous capsule, up to 3.4 mm, in contrast to 2.23 mm with unchanged joints, and an important sign was the identification of normal sizes of hyaline cartilage. At the same time, the sizes of osteophytes increased as the stage of coxarthrosis increased from 4.2 mm to 10.1 mm. Thus, in stage III coxarthrosis, the size of osteophytes was 8.3 mm, and in coxarthrosis formed as a result of aseptic necrosis of the femoral head, it increased to 15.0 mm.

All patients with stages I-III of coxarthrosis revealed significant differences in the shape of the head of the hip joint, so if at stage I of the disease almost all (96%) of the examined patients had a spherical shape of the head, then at stage II of the disease only in 58% of cases a spherical shape was detected, and in 42% of cases - a moderately flattened shape of the head. At stage III, this sign increased and in 96%

of cases it was found to be significantly flattened (Fig. 2).

Fig.2. Sonogram of the hip joint in coxarthrosis (significantly flattened shape of the head).

When studying the joint cavity in patients with stage I coxarthrosis, effusion was practically absent (1% of cases), at stage II this sign was already detected in 10% of cases, and in patients with stage III coxarthrosis, effusion was visualized in 37% of cases. That is, as the disease progresses, the likelihood of effusion in the joint cavity increases.

When applying the method of ultrasonic dopplerography, a progressive decrease in the speed indicators of blood flow in the lateral circumflex arteries of the thigh was revealed as the stage of coxarthrosis increases, so the PVR decreases from 25.2 cm/s in the first stage of coxarthrosis to 16.6 cm/s in the third stage; CDS from 4.6 cm/s at the first stage to 2.1 cm/s at the III stage of coxarthrosis. In addition, if the PSS at stage II is 22.4 cm/s, and with coxarthrosis formed as a result of aseptic necrosis of the femoral head, it significantly decreases to an average of 17.8 cm/s. In rheumatoid arthritis, there was an increase in PSS of blood flow, on average, up to 28.0 cm/s, as well as an increase in DPV of blood flow, up to 8.1 cm/s. and a decrease in IR to 0.65, when comparing the values of groups with unchanged joints and stage II of coxarthrosis at 0.78 and 0.77, respectively.

Discussion

The thickness of the hyaline cartilage and fibrous capsule, the presence or absence of intra-articular effusion, the shape of the femoral head, the size of osteophytes (if any) are the main structural parameters on which the differential diagnosis of dystrophic and inflammatory diseases of the hip joint is based. In case of an inflammatory lesion of the hip joint, the leading sonographic criteria are: an increase in PVR to numbers - 32.0 cm/s, a CDP to - 10.3 cm/s, and a decrease in IR to numbers - 0.65, combined with an increase

in the thickness of the fibrous capsules while maintaining the normal thickness of the hyaline cartilage. And the signs of the presence of coxarthrosis are: a progressive decrease in the rate of blood flow in the lateral circumflex arteries of the thigh and progressive thinning of the hyaline cartilage, the presence of marginal osteophytes, a violation of the shape and contour of the femoral head. Coxarthrosis, formed as a result of aseptic necrosis of the femoral head, is characterized by a significant decrease in the rate of blood flow in the lateral circumflex arteries of the thigh (PRS up to digits - 17.8 cm/s, CDS up to digits - 3.7 cm/s) in combination with early and more complete thinning of the hyaline cartilage, more significant sizes of osteophytes, significant deformity of the femoral head, effusion in the joint cavity.

Conclusions

A number of undoubted advantages - non-invasiveness (unlike arthroscopy), accessibility, simplicity, cost-effectiveness (compared to CT and MRI) - provided the method

of hip joint ultrasound with a priority among other instrumental methods for examining joints and soft tissues [5]. Ultrasound is highly informative in reflecting small details of the surface of bones, ligamentous tendon apparatus, and also allows you to identify and control inflammatory changes in tissues. The advantage of ultrasound over the X-ray method is safety, polypositionality, as well as the possibility of dynamic observation.

The most important features of a comprehensive ultrasound examination of the hip joints are: determination of the state of the joint capsule, the presence of effusion in the cavity of the hip joint, assessment of the sphericity and contour of the head, determination of the state of the hyaline cartilage and periarticular tissues, and using the Doppler method, the blood flow in the lateral circumflex arteries of the thigh is assessed, with the calculation of blood flow PSV, blood flow DPV and IR, which is quite important in the differentiation of inflammatory and degenerative processes.

References / Список литературы / Iqtiboslar

1. Zhang, W. EULAR recommendations for knee and hip osteoarthritis: a critique of the methodology / W. Zhang, M. Doherty // Br. J. Sports Med. - 2006. - Vol. 40. - p. 664-669.
2. Kovalenko, V.N. Osteoporosis. Practical guidance / N. Kovalenko, O.P. Bortkevich. - K.: Morion, 2003. - 448 p.
3. Khabirov, R.A. Ultrasound differential diagnosis of degenerative and inflammatory joint lesions / R.A. Khabirov, M.G. Tukhbatulin, E.R. Volkova // Nauch. - pract. rheumatology. - 2001. - No. 3. - p. 125.
4. Yaremenko, O.B. Modern algorithm for diagnosing rheumatoid arthritis / O.B. Yaremenko // Clinical Immunology. Allergology. Infectology. - Kiev, 2006. - No. 2. - p. 54-58.
5. Saltykova, V.G. Possibilities of ultrasound diagnostics in injuries and diseases of large joints / V.G. Saltykova, A.K. Morozov // Proceedings of the All-Russian Congress of Radiation Diagnostics. - M., 2007. - p. 316-317.
6. Divakov M.G., Boloboshko K.B. Early ultrasound diagnosis of osteochondropathy of the femoral head // News of Radiation Diagnostics. 1999; 2:12-13.
7. Шамсиев А.М., Хусинова С.А. (2008). Влияние факторов окружающей среды на здоровье человека в Узбекистане. В: Бенке Р. (ред.) Социально-экономические причины и последствия опустынивания в Центральной Азии. Серия «Наука НАТО для мира и безопасности». Шпрингер, Дордрехт. https://doi.org/10.1007/978-1-4020-8544-4_11
8. Хусинова Ш. А., Аблакулова М. Х. Диабетдан олдинги ҳолат //Проблемы биологии и медицины. – 2017. – №. 1. – С. 93.
9. Хусинова Ш., Аблакулова М. Лейла Хакимова Процесс рационального назначения лекарственных средств и выбор персонального лекарства в практике ВОП. – 2020.
10. Лейла Хакимова, Шоира Хусинова, Муниса Аблакулова, Дилшода Абдухамидова Безопасное материнство и эффективный антенатальный уход // ОИИ. 2021. №8/S. URL: <https://cyberleninka.ru/article/n/bezopasnoe-materinstvo-i-effektivnyu-antenatalnyu-uhod>
11. Шоира Хусинова, Муниса Аблакулова, Лейла Хакимова Процесс рационального назначения лекарственных средств и выбор персонального лекарства в практике ВОП // ОИИ. 2020. №1/S. URL: <https://cyberleninka.ru/article/n/protsess-ratsionalnogo-naznacheniya-lekarstvennyh-sredstv-i-vybor-personalnogo-lekarstva-v-praktike-vop> (дата обращения: 18.03.2023).
12. Akbarovna, K. S. (2022). UVB Therapy for the Treatment of Patients with Chronic Dermatitis. Central Asian Journal of Medical and Natural Science, 3(5), 676-678. Retrieved from <https://cajms.centralasianstudies.org/index.php/CAJMNS/article/view/1152>
13. Хусинова Ш. А., Аминов З. З. Мероприятия, проводимые в учреждениях первичной медико-санитарной помощи по профилактике распространения COVID-19 // Медицинское образование сегодня. – 2020. – №. 3. – С. 190-201.
14. Хусинова Ш. А. Распространённость артериальной гипертензии, особенности её клинического течения и состояние белково-липидного обмена у работников табачного производства // Автореф. дисс.... канд. мед. наук. Ташкент. – 1998. – С. 24.
15. Khakimova L. R., Khusinova S. A. MH Ablakulova Results of the implementation of a clinical protocol for the integrated management of patients with arterial hypertension and diabetes mellitus in primary health care. 2018 // Journal Health, demography, ecology of the Finno-Ugric peoples. – №. 4. – С. 66-68.
16. Хусинова, . Ш., Хакимова, . Л., & Курбанов, Б. (2022). РЕЗУЛЬТАТЫ ОЦЕНКИ КАЧЕСТВА ЛЕЧЕНИЯ БОЛЬНЫХ С СЕРДЕЧНОЙ НЕДОСТАТОЧНОСТЬЮ В УСЛОВИЯХ ПОЛИКЛИНИКИ. Журнал кардиореспираторных исследований, 2(3), 82–84. <https://doi.org/10.26739.2181-0974-2021-3-16>